

Response: Using Handheld Portable Ultrasound in Bedside Assessment of Post-stroke Hemiplegic Shoulders: Early Detection of Glenohumeral Subluxation

Yanıt: İnme Sonrası Hemiplejik Omuzların Yatak Başı Değerlendirilmesinde El Tipi Taşınabilir Ultrasonun Kullanılması: Glenohumeral Subluksasyonun Erken Tespiti

 Burak Tayyip DEDE^a,  Muhammed OĞUZ^b,  Bülent ALYANAK^c,  Fatih BAĞCIER^d

^aProf. Dr. Cemil Taçcıoğlu City Hospital, Clinic of Physical Medicine and Rehabilitation, İstanbul, Türkiye

^bİstanbul Training and Research Hospital, Clinic of Physical Medicine and Rehabilitation, İstanbul, Türkiye

^cKocaeli University Faculty of Medicine, Department of Physical Medicine and Rehabilitation, Kocaeli, Türkiye

^dBaşakşehir Çam and Sakura City Hospital, Clinic of Physical Medicine and Rehabilitation, İstanbul, Türkiye

Dear Editor;

We carefully read the article “Using Handheld Portable Ultrasound in Bedside Assessment of Post-stroke Hemiplegic Shoulders: Early Detection of Glenohumeral Subluxation”.¹ With the advances in technology, portable handheld ultrasound (PHU) devices have been developed and used by clinicians in many fields.^{1,2} This point is well emphasized in this article. In your article, it was stated that >2 mm difference in acromio-greater tuberosity (AGT) distance may be significant for subluxation.¹ However, it may be useful to mention a few points to be considered regarding AGT distance measurement.

Although AGT distance measurements have been reported as reliable measurements, in a study evaluat-

ing shoulder ultrasonographic measurements, the intra-rater and inter-rater minimal detectable change (MDC) values of AGT distance were 0.78 cm and 0.55 cm, respectively.³ To achieve good reliability in all age groups, they did not recommend the use of AGT distance measurement until further improvements in US protocols.⁴ The MDC value is very important and represents the smallest clinically significant difference for the measurements to be considered reliable. Furthermore, another study demonstrated the relationship between AGT distance and individual-related variables, finding a significant correlation between the individual’s height and AGT distance.⁵

Cholewinski et al. found that AGT distance was significantly different between individuals with sub-

TO CITE THIS ARTICLE:

Dede BT, Oğuz M, Alyanak B, Bağcier F. Response: Using Handheld Portable Ultrasound in Bedside Assessment of Post-Stroke Hemiplegic Shoulders: Early Detection of Glenohumeral Subluxation. Türkiye Klinikleri Journal of Physical Medicine and Rehabilitation Sciences. 2024;27(2):215-6.

Correspondence: Burak Tayyip DEDE

Prof. Dr. Cemil Taçcıoğlu City Hospital, Clinic of Physical Medicine and Rehabilitation, İstanbul, Türkiye

E-mail: drbrk22.94@gmail.com

Peer review under responsibility of Journal of Physical Medicine and Rehabilitation Science.

Received: 05 Jun 2024

Accepted: 19 Jun 2024

Available online: 09 Jul 2024

1307-7384 / Copyright © 2024 Turkey Association of Physical Medicine and Rehabilitation Specialist Physicians. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).



acromial impingement syndrome and a healthy control group without any complaints or history of shoulder pain.⁵ They reported that the difference in AGT distance between both shoulders ranged from -3.9 to +8.6 mm, with a median difference of 2.7 mm. Therefore, the reliability of the underlying cut-off value of >2 mm is questionable. In addition, Kumar and Attwood found the inter-rater standard error of measurement and MDC values to be 0.2-0.2 cm and 0.3-0.4 cm for the right and left shoulder, respectively in a reliability study of AGT distance measurement performed by experienced and novice assessors in healthy controls.³ This shows that the experience of the person performing the measurement is important.

In addition, the widespread use of PHU devices has led to questions about the reliability of these devices.

Previously, the reliability of PHU devices in measuring the vertical distance between the acromion and humerus was demonstrated.² However, to the best of our knowledge, no study has evaluated the reliability of PHU devices for AGT distance measurements.

In conclusion, more reliable measurements are needed to evaluate shoulder subluxation in patients with post-stroke hemiplegia. There is also a need for studies to evaluate the reliability of AGT distance measurements of PHU devices in clinical use.

REFERENCES

1. Erden Y, Korkut M. [Using handheld portable ultrasound in bedside assessment of post-stroke hemiplegic shoulders: early detection of glenohumeral subluxation]. *Turkiye Klinikleri Journal of Physical Medicine and Rehabilitation Sciences*. 2024;27:144-6. [[Crossref](#)]
2. Dede BT, Aytikin E, Bağcier F. Measures of acromiohumeral distance with wireless ultrasound machine in subacromial impingement syndrome: an inter-machine reliability study. *J Ultrason*. 2024;24:18. [[Link](#)]
3. Kumar P, Attwood M. Inter-rater reliability of ultrasound measurements of acromion-greater tuberosity distance between experienced and novice raters in healthy people. *Musculoskeletal Care*. 2018;16:163-6. [[Crossref](#)] [[PubMed](#)]
4. Yuan X, Lowder R, Aviles-Wetherell K, et al. Reliability of point-of-care shoulder ultrasound measurements for subacromial impingement in asymptomatic participants. *Front Rehabil Sci*. 2022;3:964613. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
5. Cholewinski JJ, Kusz DJ, Wojciechowski P, et al. Ultrasound measurement of rotator cuff thickness and acromio-humeral distance in the diagnosis of subacromial impingement syndrome of the shoulder. *Knee Surg Sports Traumatol Arthrosc*. 2008;16:408-14. [[Crossref](#)] [[PubMed](#)]