

# Central Diabetes Insipidus in a Quadriplegic Patient without Direct Trauma to the Head: A Rare Complication

## Kafa Travması Eşlik Etmeyen Kuadriplejik Bir Hastada Santral Diabetes İnsipidus: Nadir Bir Komplikasyon

<sup>1</sup> Sefa GÜMRÜK ASLAN<sup>a</sup>, <sup>2</sup> Kutay TEZEL<sup>a</sup>, <sup>3</sup> Esra ÜLGEN KIRATLIOĞLU<sup>a</sup>,  
<sup>4</sup> Nilüfer Kutay ORDU GÖKKAYA<sup>b</sup>

<sup>a</sup>University of Health Science Ankara Gaziler Physical Therapy and Rehabilitation Training and Research Hospital, Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

<sup>b</sup>Ankara Bilkent City Hospital, Clinic of Physical Medicine and Rehabilitation, Ankara, Türkiye

**ABSTRACT** This report highlights the clinical significance of central diabetes insipidus (CDI), a rare complication that arises following acute traumatic cervical spinal cord injury without head injury. The aim is to emphasize the importance of considering CDI during the rehabilitation process. A 25-year-old male patient diagnosed with C4 level (ASIA A). No head trauma was detected in the patient's examinations. The patient reported symptoms of dry mouth, excessive thirst, and increased water intake during the neurosurgery clinic evaluation. The patient received a diagnosis of CDI. During hospitalization, the patient's fluid intake ranged from 3,000-4,000 mL daily. With a strict fluid regimen and urine monitoring, the patient's laboratory findings improved within two weeks. Being aware of the potential symptoms of diabetes insipidus that may accompany spinal cord injuries enables the rapid implementation of appropriate treatment and fluid management strategies. This helps maintain the patients' overall health and minimizes potential complications.

**Keywords:** Quadriplegia; head trauma; spinal cord injury; diabetes insipidus

**ÖZET** Bu vaka, kafa travması olmaksızın akut travmatik servikal omurilik yaralanmasını takiben ortaya çıkan nadir bir komplikasyon olan santral diabetes insipidusun (SDİ) klinik önemini vurgulamaktadır. Bu çalışmanın amacı, rehabilitasyon sürecinde SDİ'nin göz önünde bulundurulmasının önemini vurgulamaktır. Yirmi beş yaşındaki erkek hasta C4 ASIA A tanısı ile takip ediliyordu. Hastanın değerlendirmelerinde kafa travmasının eşlik etmediği saptandı. Hastanın takiplerinde, ağız kuruluğu, aşırı susama ve artmış su tüketimi şikâyetleri gelişmiştir. Hastaya SDİ tanısı konulmuştur. Hastanede yatış süresince hastanın günlük sıvı alımı 3.000-4.000 mL arasında değişmiştir. Sıkı bir sıvı rejimi ve idrar takibi ile hastanın laboratuvar bulguları 2 hafta içinde düzelmiştir. Omurilik yaralanmalarına eşlik edebilecek diabetes insipidus semptomlarının farkında olunması, uygun tedavi ve sıvı yönetim stratejilerinin hızla uygulanmasını sağlar. Bu durum, hastaların genel sağlığının korunmasına yardımcı olur ve potansiyel komplikasyonları en aza indirir.

**Anahtar Kelimeler:** Kuadripleji; kafa travması; spinal kord hasarı; diabetes insipidus

Spinal cord injuries (SCI) typically result in sensory, motor, and autonomic deficits below the level of the lesion.<sup>1</sup> SCI is a traumatic event with limited recovery of function despite the best efforts of many researchers to devise realistic therapeutic

treatments. SCI primarily affects young adults, with nearly half of all injuries occurred between the ages of 16 and 30. The average age at injury has increased from 29 years during the 1970s to 43 since 2015.<sup>2,3</sup>

**TO CITE THIS ARTICLE:**

Yazar Adı. Makale Başlığı. Türkiye Klinikleri Journal of Physical Medicine and Rehabilitation Sciences. 2024;?(?):???-???

**Correspondence:** Sefa GÜMRÜK ASLAN

University of Health Science Ankara Gaziler Physical Therapy and Rehabilitation Training and Research Hospital,  
Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

**E-mail:** sefagumrukk@gmail.com



Peer review under responsibility of Journal of Physical Medicine and Rehabilitation Science.

**Received:** 26 Jun 2024

**Received in revised form:** 15 Oct 2024

**Accepted:** 24 Oct 2024

**Available online:** 05 Nov 2024

1307-7384 / Copyright © 2024 Turkey Association of Physical Medicine and Rehabilitation Specialist Physicians. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

The National Spinal Cord Injury Statistics Center data is estimated that the annual of SCI incidence is approximately 54 cases per million people, or 17,810 new cases each year in the U.S.<sup>2</sup>

Published data about the epidemiology of SCI in Türkiye is limited to a few studies. Karacan et al. reported an epidemiological study conducted by 49 provinces, and accordingly states that the annual of SCI incidence is 12.7 cases per million people.<sup>4</sup>

Since 2015, motor vehicle crashes take the first place of reported SCI cases (38.6%). The most common causes include falls (32.2%), followed by acts of violence (primarily gunshot wounds, 14%) and recreational sports activities (7.8%).<sup>2</sup> Violence caused 13.3% of SCI prior to 1980, and peaked between 1990 and 1999 at 24.8% before declining to only 14.3% since 2010.<sup>3</sup> In Türkiye, the most common cause of injury is motor vehicle accidents (48.8%), following by falls (36.5%), stab wounds (3.3%), violence (1.9%), and injuries after diving (1.2%).<sup>4</sup>

The most common levels of SCI injury were C5 and C6 among tetraplegics and T12 and L1 among paraplegics. C5 is the most common vertebral damage level, followed by C4, C6, T12, and L1 levels.<sup>5</sup> Recently, incomplete tetraplegia is the most frequent neurological category at discharge of persons reported the database (47.2%), followed by incomplete paraplegia (19.6%), complete paraplegia (20.2%) and complete tetraplegia (12.3%). Less than 1% of persons experienced complete neurological recovery by the time of hospital discharge.<sup>2,5</sup>

Secondary complications were present during both acute care units and inpatient rehabilitation services. These include pulmonary, cardiovascular, gastrointestinal complications, bladder and bowel problems, spasticity, pressure ulcers, neuropathic pain, syringomyelia, osteoporosis, autonomic dysreflexia, and upper extremity overuse syndromes. The presence of secondary complications may decrease health-related quality of life, limit function recovery and increase in morbidity and mortality rates.<sup>6</sup>

Diabetes insipidus is a rare disorder of water homeostasis characterized by the excretion of excessive volumes of hypotonic urine (polyuria). Over 90% of the vasopressinergic neurons which project

from the supraoptic and paraventricular nuclei to terminate in the posterior pituitary must be destroyed in order to cause vasopressin (AVP) deficiency sufficient to lead polyuric symptoms. Central diabetes insipidus (CDI) is the clinical manifestation of destruction of the neurons of the hypothalamus/posterior pituitary axis, with consequent loss of AVP secretion. In many patients, CDI is caused by the destruction or degeneration of neurons originating in the supraoptic and paraventricular nuclei. It is clinically characterised by polydipsia and polyuria (urine output >30 mL/kg/day) of dilute urine (<250 mOsm/L).<sup>7</sup>

Diabetes insipidus arises from one of several causes and can be classified as either primary or secondary.<sup>8</sup> The primary causes are either idiopathic or familial. The secondary group consists mainly of local inflammatory or autoimmune diseases, cranio-cerebral traumas, Langerhans cell histiocytosis, infections, neoplastic diseases, trauma resulting from surgery or an accident (e.g SCI), metastases and midline cerebral and cranial malformations.<sup>9-11</sup> In the presence of polyuria and polydipsia symptoms developing during the follow-up of patients with spinal cord injury, the diagnosis of CDI should not be overlooked.<sup>12,13</sup>

We present a case of CDI that developed after a gunshot injury to the cervical spinal cord unassociated with head injury. We aimed to discuss the disease process in light of the relevant literature and to emphasize restoring and compensating for the loss of functioning, and preventing or slowing deterioration in functioning in every area of a person's life during the rehabilitation process.

## CASE REPORT

A 25-year-old male patient applied to the emergency department following a gunshot wound to his posterior neck region. The computer tomography (CT) taken in the emergency room, it was found that the bullet core was placed in the cervical (C5) vertebra corpus and caused a fracture in the left lamina of the C6 vertebra. In cervical spinal magnetic resonance imaging (MRI), heterogeneous signal changes were observed in the posterior elements at the level of C5-

C6 vertebrae, and heterogeneous signal changes were detected in an area of approximately 13x24 mm in the cervical spinal cord at this level. No mass lesion that can be differentiated from the brain parenchyma was detected in brain CT. The cerebral parenchyma, posterior fossa, and cerebral sulcus were normal and the ventricles were symmetrical and in normal calibration. In addition, the midline structures are in place and no lytic-destructive lesion was observed in the bone structures. Radiographs of the thoracic spine showed no vertebral abnormalities. The abdominal CT revealed no acute pathology. The patient was operated by the neurosurgery clinic and the bullet in the C5 was removed, and unilateral laminectomy was performed in the left C6 vertebra. The patient is not indicated for a vertebral stabilization operation.

The patient, who was taken to the physical therapy and rehabilitation program in the early postoperative period, was transferred to our hospital for an advanced rehabilitation program in the fourth week after his medical condition stabilized. At initial physical examination, the general condition of the patient was good, he was conscious, his cooperation and orientation were complete. The bilateral shoulder elevation muscle strength were 5/5, shoulder abduction 2/2, elbow flexion (C5) 3/1 +, wrist extensors (C6) and other distal muscles were 0/5 based on the Global Oxford Scale. The deep tendon reflexes of the upper and lower extremities were hypoactive. On sensory examinations; C5 and distal dermatomes were anesthetic. Pathological reflexes were negative. No sensory nor motor functions are preserved in the sacral segments S4-5. The patient was at the in-bed level and there was no support/unsupported sitting balance. The patient's neurological examinations were determined C4 ASIA A with sensory and motor examination without deep anal pressure sensation and internal anal sphincter contraction.

In the history of the patient has been learned, he complained of dry mouth, drink too much water and excessive thirst on the neurosurgery clinic. After the endocrine and psychiatric consultations and examinations requested for differential diagnosis the patient was diagnosed with central diabetes insipidus and initiated orally desmopressin acetate 120 mcg/day.

In early period of his rehabilitation program includes passive range of motion exercises for the lower and upper extremities twice a day, progressive resistive and active assistive exercises to increase existing muscle strength, breathing exercises, vascular endurance program on the gradual tilt table and mobilization programs to increase trunk balance. During the follow-up of the patient, occasional hypotension attacks were detected. The hemodynamic status of the patient was followed closely.

During the hospitalization follow-up, it was observed that the daily fluid intake was around 3,000-4,000 mL, and the urine output varied between 6,400 and 7,400 mL and continued at a very high level. Laboratory examinations of the patient after the introduction and drug regulation are given in Table 1. Blood pressure values were found to be within normal limits. Desmopressin acetate dose of the patient was gradually increased up to 240 mcg/day in consultation with internal medicine and nephrology during the decision-making phase in terms of monitoring the patient's current polyuria and fluid deficit. We monitored the patient's urea and electrolytes regularly, for evidence of dehydration, renal hypoperfusion, or electrolyte imbalance. Laboratory findings improved after two weeks of strict fluid regimen and urine follow-up. It was decided to continue the desmopressin acetate dose as 240 mcg/day. At follow-up all his laboratory parameters were normal. Our patient was still using a dose of 240 mcg/day in follow-up at the 3<sup>rd</sup> month. The participant provided written informed consent.

**TABLE 1:** Laboratory results before and after treatment.

	Before treatment	After treatment
Serum sodium (mEq/L)	134	138
GFR (mL/minute/1.73 m <sup>2</sup> )	162	145
Creatinin (mg/dL)	0.42	0.55
Chloride (mmol/L)	100	103
Ca (mg/dL)	9.9	9.5
Urine specific gravity (N.1020-1028)	1.010	1020-1030
Urine osmolality (mOsm/L)	143	184
Serum osmolality (mOsm/L)	284	285
Daily urine output	6400-7400	2250-3000

GFR: Glomerular filtration rate; Ca: Calcium.

## DISCUSSION

Head injury and SCI are common occurrences in the neuro-surgical practice. They account for an important proportion of care, reduce disability, and ethical issues. The reported incidence of cervical spine trauma after clinically significant head injury generally ranges from 4% to 8%.<sup>14</sup> Hypercalcemia is common in children (24%) and young adults after SCI. Hypercalciuria develops within the first week after injury and can last 6 months or longer. Symptoms typically present 4 to 8 weeks after SCI. Symptoms include anorexia, nausea, lethargy, polydipsia, and polyuria. As early as three months, bones become osteoporotic and long bone fractures can be seen. It has been shown that parathyroid hormone and vitamin D levels are significantly suppressed in patients after SCI. It has been reported that this suppression returns to normal within six months.<sup>14</sup> As the patient's blood calcium levels, as well as vitamin D levels, were within normal ranges upon admission, it was concluded that the patient's symptoms were unlikely to be attributable to hypercalcemia.

Berezin et al. reported six women with a traumatic SCI developed hyperprolactinemia, amenorrhea and galactorrhea. Five of them had thoracic level lesions and 1 had a lumbosacral lesion. One of the cases developed transient diabetes insipidus. The authors hypothesizes one or more of the following factors might play a role in the pathogenesis of this syndrome: pituitary stalk concussion or contusion, the elevated level of endorphins lasted only for the duration of the spinal shock, or injury of the afferent neuronal pathway of the spinal cord.<sup>15</sup>

The initial mechanical in acute SCI forces delivered to the spinal cord at the time of injury is known as primary injury where "displaced bone fragments, disc materials, and/or ligaments bruise or tear into the spinal cord tissue".<sup>16</sup> Four main characteristic mechanisms of primary injury have been identified that include: (1) Impact plus persistent compression; (2) Impact alone with transient compression; (3) Distraction; (4) Laceration/transection.<sup>17</sup> Damage in this process is irreversible.

Secondary injury can be temporally divided into acute, sub-acute, and chronic phases. The acute phase

begins immediately following SCI and includes vascular damage, ionic imbalance, neurotransmitter accumulation (excitotoxicity), free radical formation, calcium influx, lipid peroxidation, inflammation, edema, and necrotic cell apoptosis. As the injury progresses, the sub-acute phase of injury begins which involves apoptosis, demyelination of surviving axons, Wallerian degeneration, axonal dieback, matrix remodeling, and evolution of a glial scar around the injury site. Further changes occur in the chronic phase of injury including the formation of a cystic cavity, progressive axonal die-back, and maturation of the glial scar.<sup>18-20</sup>

In addition to direct tissue damage due to gunshot wound injuries, permanent tissue damage such as ischemia, thrombosis, and necrosis should be kept in mind due to thermal effects as well as the direct impact of the bullet, the pressure of shock waves and the temporary cavitation effect in surrounding tissues, especially in neural structures. Three mechanisms of tissue damage: the direct impact of the bullet, the pressure of shock waves and the temporary cavitation after gunshot wounds injuries.<sup>21</sup> Mirovsky et al. verified this, cause of neurologic deficit in patients following gunshot wounds is direct trauma to the spinal cord by the bullet, bone fragments, or in rare cases by disc fragments compressing the cord.<sup>22</sup>

Some of the underlying causes of CDI are iatrogenic post neurosurgery (20%), head trauma (16%) and inherited/familial (1%).<sup>16</sup> Posttraumatic CDI can be caused by any type of head trauma however most commonly occurs secondary to road traffic accidents.<sup>23</sup> Trauma can cause anterior pituitary dysfunction or CDI and often occurs in patients with associated skull fractures or neurological defects.<sup>24,25</sup> In a small number of cases (6%) no CT/MRI finding is demonstrated, and hypoxic damage or diffuse axonal injury is presumed to be the cause.<sup>26</sup>

Machiedo et al. reported the development of CDI in three cases following with non-cranial trauma are presented. They associated the etiology of CDI in patients, the mechanism that best explains the syndrome is shock from hypovolemia with occlusion of vessels resulting in tissue hypoxia and necrosis.<sup>27</sup>



Hypotension is commonly observed during the spinal shock phase following a spinal cord injury, primarily due to reduced sympathetic tone and vasodilation. CDI can rarely occur in adults without trauma as a consequence of hypoxemia and hypoperfusion.<sup>28</sup>

Traumatic SCI in all spine segments occurs in 5% to 15% of patients with severe head injury and may be accompanied by head trauma complications. Although an increasing injury severity has been associated, as measured by the Glasgow Coma Scale, 7 severe and moderate head injury may cause misdiagnosis of SCI leading to devastating long-term consequences.<sup>29,30</sup> Should be alert to the possibility of diabetes insipidus that may develop after SCI and fluid management is a critical aspect of SCI patient care, especially in the following CDI inpatient medical setting. Fluid management is important and vital in that patients and must be careful consideration of their individual fluid needs.

Especially in acute rehabilitation clinics, the possibility of CDI should be kept in mind by taking into account the symptoms such as polydipsia and polyuria, and by performing strict and careful fluid

intake and urine discharge follow-ups along with routine laboratory analyses. Therefore, mortality can be prevented by early diagnosis and treatment of life-threatening complications such as CDI.

#### **Source of Finance**

*During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.*

#### **Conflict of Interest**

*No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.*

#### **Authorship Contributions**

**Idea/Concept:** Sefa Gümruk Aslan; **Design:** Sefa Gümruk Aslan; **Data Collection and/or Processing:** Sefa Gümruk Aslan, Esra Ülgen Kiratlıoğlu; **Analysis and/or Interpretation:** Sefa Gümruk Aslan, Esra Ülgen Kiratlıoğlu, Kutay Tezel; **Literature Review:** Sefa Gümruk Aslan, Esra Ülgen Kiratlıoğlu, Kutay Tezel; **Critical Review:** Nilüfer Kutay Ordu Gökkaya.

## REFERENCES

1. Kirshblum SC, Burns SP, Biering-Sorensen F, et al. International standards for neurological classification of spinal cord injury (revised 2011). *J Spinal Cord Med.* 2011;34:535-46. PMID: 22330108; PMCID: PMC3232636.
2. Jain NB, Ayers GD, Peterson EN, Harris MB, Morse L, O'Connor KC, Garshick E. Traumatic spinal cord injury in the United States, 1993-2012. *JAMA.* 2015 Jun 9;313(22):2236-43. PMID: 26057284; PMCID: PMC4712685.
3. Jackson AB, Dijkers M, Devivo MJ, et al. A demographic profile of new traumatic spinal cord injuries: change and stability over 30 years. *Arch Phys Med Rehabil.* 2004;85:1740-8. PMID: 15520968.
4. Karacan I, Koyuncu H, Pekel O, et al. Traumatic spinal cord injuries in Turkey: a nation-wide epidemiological study. *Spinal Cord.* 2000;38:697-701. PMID: 11114778.
5. Devivo MJ. Epidemiology of traumatic spinal cord injury: trends and future implications. *Spinal Cord.* 2012;50:365-72. PMID: 22270188.
6. Wahman K, Nilsson Wikmar L, Chlaidze G, et al. Secondary medical complications after traumatic spinal cord injury in Stockholm, Sweden: Towards developing prevention strategies. *J Rehabil Med.* 2019;51:513-7. PMID: 31243468.
7. Gubbi S, Hannah-Shmouni F, Koch CA, et al. Diagnostic Testing for Diabetes Insipidus. 2022 Nov 28. In: Feingold KR, Anawalt B, Blackman MR, et al, editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. PMID: 30779536.
8. Hui C, Khan M, Khan Suheb MZ, et al. Arginine Vasopressin Disorder (Diabetes Insipidus). 2024. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. PMID: 29262153.
9. Alaca R, Yilmaz B, Gunduz S. Anterior hypopituitarism with unusual delayed onset of diabetes insipidus after penetrating head injury. *Am J Phys Med Rehabil.* 2002;81:788-91. PMID: 12362120.
10. Bode H, Crawford J, Danon M. Disorders of antidiuretic hormone homeostasis: diabetes insipidus and SIADH. In: Lifshitz F, ed. *Pediatric Endocrinology*. 1st ed. New York: Marcel Dekker; 1996. p.731-52.
11. Maghnie M, Cosi G, Genovese E, et al. Central diabetes insipidus in children and young adults. *N Engl J Med.* 2000;343:998-1007. PMID: 11018166.
12. Closson BL, Beck LA, Swift MA. Diabetes insipidus and spinal cord injury: a challenging combination. *Rehabil Nurs.* 1993;18:368-74. PMID: 7938892.
13. Farrell CA, Staas WE Jr. Diabetes insipidus in a quadriplegic patient. *Arch Phys Med Rehabil.* 1986;67:132-4. PMID: 3954565.
14. McKinley WO, Gittler MS, Kirshblum SC, et al. Spinal cord injury medicine. 2. Medical complications after spinal cord injury: Identification and management. *Arch Phys Med Rehabil.* 2002;83:S58-64, S90-8. PMID: 11973698.
15. Berezin M, Ohry A, Shemesh Y, et al. Hyperprolactinemia, galactorrhea and amenorrhea in women with a spinal cord injury. *Gynecol Endocrinol.* 1989;3:159-63. PMID: 2510463.

- 
16. Sekhon LH, Fehlings MG. Epidemiology, demographics, and pathophysiology of acute spinal cord injury. *Spine (Phila Pa 1976)*. 2001;26:S2-12. PMID: 11805601.
  17. Rowland JW, Hawryluk GW, Kwon B, et al. Current status of acute spinal cord injury pathophysiology and emerging therapies: promise on the horizon. *Neurosurg Focus*. 2008;25:E2. PMID: 18980476.
  18. Oyinbo CA. Secondary injury mechanisms in traumatic spinal cord injury: a nugget of this multiply cascade. *Acta Neurobiol Exp (Wars)*. 2011;71:281-99. PMID: 21731081.
  19. Tran AP, Warren PM, Silver J. The biology of regeneration failure and success after spinal cord injury. *Physiol Rev*. 2018;98:881-917. PMID: 29513146; PMCID: PMC5966716.
  20. Dyck SM, Karimi-Abdolrezaee S. Chondroitin sulfate proteoglycans: key modulators in the developing and pathologic central nervous system. *Exp Neurol*. 2015;269:169-87. PMID: 25900055.
  21. Sidhu GS, Ghag A, Prokuski V, et al. Civilian gunshot injuries of the spinal cord: a systematic review of the current literature. *Clin Orthop Relat Res*. 2013;471:3945-55. PMID: 23479233; PMCID: PMC3825909.
  22. Mirovsky Y, Shalmon E, Blankstein A, et al. Complete paraplegia following gunshot injury without direct trauma to the cord. *Spine (Phila Pa 1976)*. 2005;30:2436-8. PMID: 16261122.
  23. Capatina C, Paluzzi A, Mitchell R, et al. Diabetes Insipidus after Traumatic Brain Injury. *J Clin Med*. 2015;4:1448-62. PMID: 26239685; PMCID: PMC4519799.
  24. Sandean D. Management of acute spinal cord injury: a summary of the evidence pertaining to the acute management, operative and non-operative management. *World J Orthop*. 2020;11:573-83. PMID: 33362993; PMCID: PMC7745491.
  25. Christ-Crain M. EJE AWARD 2019: new diagnostic approaches for patients with polyuria polydipsia syndrome. *Eur J Endocrinol*. 2019;181:R11-R21. PMID: 31067508; PMCID: PMC6598864.
  26. Adams NC, Farrell TP, O'Shea A, et al. Neuroimaging of central diabetes insipidus-when, how and findings. *Neuroradiology*. 2018;60:995-1012. PMID: 30097693.
  27. Machiedo G, Bolanowski PJ, Bauer J, et al. Diabetes insipidus secondary to penetrating thoracic trauma. *Ann Surg*. 1975;181:31-4. PMID: 1119865; PMCID: PMC1343710.
  28. Wilson RH, Whiteside MC, Moorehead RJ. Problems in diagnosis and management of hypovolaemia in spinal injury. *Br J Clin Pract*. 1993;47:224-5. PMID: 8260349.
  29. Holly LT, Kelly DF, Counelis GJ, et al. Cervical spine trauma associated with moderate and severe head injury: incidence, risk factors, and injury characteristics. *J Neurosurg*. 2002;96:285-91. PMID: 11990836.
  30. Kiboi J, Omar A, Omar MA. Head injury with concurrent cervical spine injury. *Annals of African Surgery*. 2019;16. doi: 10.4314/aas.v16i1.3.