

Dysphagia Management in Acute Stroke Patients: The Result of a Prospective Close Follow-Up Design

Akut İnme Hastalarında Disfaji Yönetimi: Prospektif Yakın Takip Tasarımının Sonuçları

^aNazife KAPAN^a, ^bMüessers ARAS^b, ^cEmre ADIGÜZEL^{b,c}, ^dMerve KAPAN^d, ^eMariam KAVAKCI^e,
^fMelike URAL^d, ^gRojda ÖZALP^d, ^hSeher Merve TİRYAKİ^d, ⁱHesna BEKTAŞ^f, ^jEvren YAŞAR^g

^aKırşehir Ahi Evran University Faculty of Medicine, Department of Physical Medicine and Rehabilitation, Kırşehir, Türkiye

^bAnkara Yıldırım Beyazıt University Faculty of Medicine, Ankara Bilkent City Hospital, Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

^cUniversity of Health Sciences Gülhane Faculty of Medicine, Health Application and Research Center,
Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

^dAnkara Bilkent City Hospital, Clinic of Physical Medicine and Rehabilitation, Ankara, Türkiye

^eAnkara Yıldırım Beyazıt University Faculty of Medicine, Ankara Bilkent City Hospital, Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

^fAnkara Bilkent City Hospital, Clinic of Neurology, Ankara, Türkiye

^gUniversity of Health Sciences Gülhane Faculty of Medicine, Health Application and Research Center,
Department of Physical Medicine and Rehabilitation, Ankara, Türkiye

This study was presented as a poster at the VIII. Medical Rehabilitation Congress, November 9-13, 2022, Ankara, Türkiye.

ABSTRACT Objective: This study was conducted to examine the results of early rehabilitation and follow-up procedures in terms of dysphagia in acute stroke patients. **Material and Methods:** A total of 55 patients aged 18-85 years who were diagnosed with acute stroke between June 2020 and January 2021 and who were detected to have dysphagia by bedside swallowing test were included for prospective follow-up. Patients who were unconscious and uncooperative, had oropharyngeal structural damage or head/neck malignancy were excluded from the study. Gugging Swallow Screening Test and Eating Assessment Tool-10 were used for swallowing evaluations of the patients. Swallowing evaluation was performed before rehabilitation, at discharge and after rehabilitation. Patients were given individual early rehabilitation consisting of diet modification, safe swallowing recommendations, oral hygiene and exercise program and acute stroke patients were closely followed up during their hospital stay and for 1 month after discharge. **Results:** A significant improvement was found in dysphagia scores after the rehabilitation ($p<0.05$). A significant relationship was found between dysphagia score change and oral hygiene, stroke history, and stroke type after the rehabilitation ($p<0.05$). A negative correlation was found between the dysphagia score and the National Institutes of Health Stroke Scale score before the rehabilitation, and a positive correlation was found between the body mass index, Barthel and Functional Ambulation Classification score ($p<0.05$). **Conclusion:** In this study, it was observed that close monitoring of acute stroke patients with dysphagia with individual rehabilitation improved dysphagia scores and reduced morbidities that may develop in patients.

Keywords: Dysphagia; stroke; dysphagia rehabilitation

ÖZET Amaç: Bu çalışma, akut inmeli hastalarda disfaji açısından erken rehabilitasyon ve takip işlemlerinin sonuçlarını incelemek amacıyla yapıldı. **Gereç ve Yöntemler:** Haziran 2020 ile Ocak 2021 tarihleri arasında akut inme tanısı konulan ve yatak başı yutma testi ile disfaji tespit edilen 18-85 yaş arası 55 hasta prospektif takip için çalışmaya dahil edildi. Bilinci kapalı ve koopere olmayan, orofaringeal yapısal hasarı veya baş/boyun malignitesi olan hastalar çalışmadan hariç tutuldu. Hastaların yutma değerlendirmeleri için Gugging Yutma Tarama Testi ve Yeme Değerlendirme Aracı-10 kullanıldı. Yutma değerlendirmesi rehabilitasyondan önce, taburcu olurken ve rehabilitasyondan sonra yapıldı. Hastalara yutma değerlendirmesi sonrası diyet modifikasyonu, güvenli yutma önerileri, oral hijyen ve egzersiz programından oluşan bireysel erken rehabilitasyon uygulanarak akut inmeli hastalar hastanede yatış süreleri ve taburcu olduktan sonra da 1 ay süreyle yakın takip edilmiştir. **Bulgular:** Rehabilitasyondan sonra disfaji skorlarında anlamlı iyileşme bulundu ($p<0,05$). Rehabilitasyondan sonra disfaji skoru değişimi ile ağız hijyeni, inme öyküsü ve inme tipi arasında anlamlı ilişki bulundu ($p<0,05$). Rehabilitasyon öncesi disfaji skoru ile Ulusal Sağlık Enstitüleri İnme Ölçeği skoru arasında negatif korelasyon, beden kitle indeksi, Barthel ve Fonksiyonel Ambulasyon Sınıflandırma skoru arasında ise pozitif korelasyon bulundu ($p<0,05$). **Sonuç:** Bu çalışmada disfaji saptanan akut inme hastalarının bireysel rehabilitasyon ile yakından takip edilmesinin disfaji skorlarını iyileştirdiği ve hastalarda gelişebilecek morbiditeleri azalttığı gözlemlendi.

Anahtar Kelimeler: Disfaji; inme; disfaji rehabilitasyonu

Correspondence: Nazife KAPAN

Kırşehir Ahi Evran University Faculty of Medicine, Department of Physical Medicine and Rehabilitation, Kırşehir, Türkiye

E-mail: nazifekapan@gmail.com



Peer review under responsibility of Journal of Physical Medicine and Rehabilitation Science.

Received: 23 Sep 2024

Received in revised form: 12 Dec 2024

Accepted: 13 Dec 2024

Available online: 18 Dec 2024

1307-7384 / Copyright © 2024 Turkey Association of Physical Medicine and Rehabilitation Specialist Physicians. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Dysphagia occurs as a result of disturbances in the oral, pharyngeal, or esophageal phases of swallowing. Food aspiration after dysphagia is a strong risk factor for pneumonia and requires nutritional intervention. Therefore, oropharyngeal dysphagia may impair nutrition, hydration, and pulmonary status of the individual and may lead to mortality, especially in elderly patients. Management of pulmonary complications caused by aspiration is difficult. Evaluation of the presence and severity of aspiration occurring during the pharyngeal phase of swallowing is of primary importance for prognosis and rehabilitation. Since oropharyngeal dysphagia is often present after acute stroke and may persist in many patients, swallowing function should be evaluated as early as possible in all stroke patients.^{1,2}

Guidelines recommend early screening of stroke patients for dysphagia prior to oral intake, and there is evidence that screening practices can prevent pneumonia.^{3,4} The Stroke Care Routine Handbook, published in Brazil in 2013, recommends that all stroke patients who can maintain postural control and stay alert for 15 minutes should be screened for dysphagia and that oral feeding should be allowed only after this assessment. However, it was emphasized that comprehensive screening cannot be performed in many hospitals due to the lack of professionals specialized in this field.⁵

Early diagnosis and implementation of an appropriate rehabilitation protocol reduces the effects of dysphagia and the associated mortality. The aim of dysphagia rehabilitation is to prevent aspiration, dehydration, and malnutrition by restoring functional and physiological swallowing. There is no standard management for dysphagia rehabilitation in the literature.⁶

This study aimed to establish a management in the diagnosis and treatment process of dysphagia in acute stroke patients and to determine the factors affecting the severity and prognosis of dysphagia.

MATERIAL AND METHODS

STUDY DESIGN AND PARTICIPANTS

This clinical study was conducted as a prospective controlled study. A total of 55 patients, 26 females

and 29 males, aged 18-85 years, diagnosed with acute stroke in the Neurology Clinic between June 2020 and January 2021, and in whom dysphagia was detected by a bedside swallow test were included for prospective follow-up. Patients who were unconscious and uncooperative, and who had oropharyngeal structural damage or head/neck malignancy were excluded from the study.

The patients' demographic information and histories were obtained and a neurological examination was performed. Clinical parameters were evaluated with the National Institutes of Health Stroke Scale (NIHSS), the Barthel Index (BI), and the Functional Ambulation Classification (FAC). Swallowing assessment of the patients was performed with the Gugging Swallowing Screen (GUSS) test and the Eating Assessment Tool-10 (EAT-10) 3 times, namely, before the rehabilitation (T1), at the discharge (T2), and after the rehabilitation (T3). According to the results of the GUSS test, suitable patients with moderate and severe dysphagia were determined by videofluoroscopy (VFS) and Penetration-Aspiration Scale (PAS) before the rehabilitation program. Patients who underwent an early rehabilitation program were followed for mortality and morbidity during hospitalization and for 1 month after discharge.

The study meets the guidelines of the Declaration of Helsinki and was approved by Yıldırım Beyazıt University Faculty of Medicine Clinical Research Ethics Committee (date: June 10, 2020, no: 41). All subjects provided written informed consent. Informed consent was obtained from the parents/legal guardians of the illiterates involved in study.

DYSPHAGIA ASSESSMENT

The GUSS test consists of two main parts. The first part is called the preliminary or indirect swallow test. Patients who get 5 points from this section, which is based on observation without any application, can move on to the second section. The second part is called the direct swallow test. In this section, semi-solid (1/3, 1/2, 3-5 teaspoons thickened water-pudding consistency), liquid (3, 5, 10, 20, 50 ml water), and solid (a piece of bread) consistencies are given and evaluated. A maximum of 5 points is taken from each consistency and a total of 20 points is reached.

Scores obtained from the test are classified as follows: 0-9 points=severe dysphagia and high risk of aspiration, 10-14 points=moderate dysphagia and moderate risk of aspiration, 15-19 points=mild dysphagia and mild aspiration, and 20 points=normal. If a score of less than 5 is obtained in any of the sections, it is recommended to stop the test and perform further evaluation (VFS or fiberoptic endoscopic evaluation of swallowing). In order for this test to be applied, 24 hours must have passed since the stroke and the patient must be cooperative and be able to sit upright in bed. Its advantages over other bedside screening tests are as follows: it can evaluate all consistencies separately and regulate the patient's diet according to the degree of swallowing disorder. Its Turkish validity and reliability study was performed by Umay et al. in 2018.⁷

The EAT-10 is a 10-item scale used to evaluate dysphagia symptoms, severity, and response to treatment. Each item is scored from 0 to 4. Zero indicates no problem, and 4 indicates severe problem. The total score ranges from 0 to 40. A score of 3 or more is considered abnormal.⁸ Its Turkish validity and reliability study was performed by Demir et al. in 2016.⁹

The PAS provides information on the presence and severity of penetration and aspiration. It is scored between 1 and 8 (1=no penetration and aspiration, 2-5=penetration available, 6-8=aspiration available).¹⁰

CLINICAL PARAMETERS

The NIHSS is a scale that examines neurological functions in stroke patients and provides insight into long-term prognosis. It is a neurological assessment scale with 11 categories that provides information on consciousness level, motor function, sensory function, and speech and language functions, where 0-6 points indicate good prognosis, 7-15 points indicate moderate prognosis, and 16-42 points indicate poor prognosis.¹¹

The BI, which is used to measure the level of disability experienced by the patient during activities of daily living, consists of a total of 10 main items. The total score is evaluated between 0 and 100, where 0-20 points: fully dependent, 21-61 points: severely dependent, 62-90 points: moderately dependent, 91-99: mildly dependent, and 100 points: fully independent.¹²

FAC is used to determine the ambulation levels of patients. In this classification, ambulation of patients is evaluated at a total of 6 levels between 0 and 5.¹³

REHABILITATION PROTOCOL

After clinical and swallowing evaluation of acute stroke patients with dysphagia, an individual early rehabilitation protocol consisting of diet modification, safe swallowing recommendations, oral hygiene and exercise program determined by the authors was applied to all patients.

According to the clinical condition of the patients, a diet was recommended using the International Dysphagia Diet Standardization Initiative (IDDSI). There are 8 different levels (0-7) for food structure and liquid consistency in IDDSI. These levels are designed to be used in dysphagia patients of all age groups and in different cultures. Simple measurement methods have been used to test nutrient composition or liquid viscosity: a fork test is used for foods, and a gravity flow test from a 10 ml syringe is used for liquids.¹⁴

Oral hygiene practices were explained in detail by informing the patient and their relatives about oral care and its importance. Due to the inadequate functional status of stroke patients in the acute period, oral care was performed by their relatives in most patients.

The exercise program was designed as twice a day in 3 sets with 15 repetitions in each set, individually for each patient, according to the clinical evaluation results of the patients. The exercise program included effortful swallowing, posture, thermal tactile stimulation from oral sensory motor development techniques, lip, jaw joint, tongue range of motion and resistant tongue base exercises. The exercise program was shown in Figure 1.

It was explained with practice by a speech and language therapist to the patient and their relatives. Exercises were performed under the supervision of a therapist 2 days a week, and as a home exercise program on other days and for 1 month after discharge.

STATISTICAL ANALYSIS

Statistical analyses were performed using the IBM SPSS 23.0 package (USA) program. Descriptive data




FIGURE 1: Examples from the exercise program.

are expressed as mean, standard deviation, minimum, maximum, median, frequency, and percentage. Conformity of continuous numerical variables to normal distribution was checked with the Shapiro-Wilk test. The independent samples t-test was used for the comparison of the variables satisfying the assumption of normal distribution between paired groups, and the One-way ANOVA test was used for the comparison between more than two groups. The Mann-Whitney U test was used to compare variables that did not fit normal distribution between groups, and the Kruskal-Wallis test was used for comparisons between more than two groups. In numerical variables that were significantly different between more than two groups, the group that caused the difference was determined via Bonferroni correction. The Freidman test was used to compare three consecutive numerical measurements, and the Wilcoxon signed-rank test was used for pairwise comparisons. Pearson's chi-square and Fisher exact tests were used to compare categorical variables between groups. McNemar's test was used to compare consecutive categorical measures. Spearman or Pearson correlation tests were used to evaluate the relationships between the measurements, according to the assumption of normal distribution. If the correlation coefficient (r) was close to 1, it was interpreted as a stronger correlation, and if it was close to 0, it was interpreted as a weaker correlation. The results were obtained at the 95% confidence interval and at the $p < 0.05$ significance level.

Power analysis was performed before the study, and for alpha (H_0): 0.05 and power ($1-\beta$) (H_1): 0.95, the effect size was calculated as 0.5. The appropriate sample size was calculated as 45 based on these data. Considering the possibility of drop-out in the controls, 55 patients were included in the study.

RESULTS

DESCRIPTIVE CHARACTERISTICS OF THE PATIENTS

The sociodemographic, comorbidity, and stroke characteristics of the patients are shown in Table 1. The mean age of the patients was 65.31 ± 13.65 years, and the mean body mass index (BMI) was 28.8 ± 5.33 kg/m².

DISTRIBUTION OF CLINICAL PARAMETERS

According to the GUSS score, 25.5% of the patients in T1, 23.6% in T2, and 3.6% in T3 had severe dysphagia.

According to the GUSS result, in patients with moderate and severe dysphagia ($n=36$) who were clinically stable, who had sitting balance, who had no contrast material allergy, who could initiate swallowing, and who had a negative coronavirus disease-2019 PCR test ($n=20$), PAS values were recorded by performing VFS. The flow diagram of patients who underwent VFS is shown in Figure 2.

It was determined that the GUSS T1 score did not change statistically in T2 ($p=1.0$), but increased in T3 at a statistically significant level ($p < 0.001$).

TABLE 1: Distribution of patients' sociodemographic, comorbidity, and stroke characteristics.

Variables		Number (n)	Percentage (%)
Sex	Female	26	47.3
	Male	29	52.7
Activity	Mild activities	5	9.1
	Moderate activities	34	61.8
	High-intensity activities	16	29.1
Smoking status	Yes	18	32.7
	No	37	67.3
Alcohol use	Yes	3	5.5
	No	52	94.5
Comorbidity	HT	21	38.2
	HT-DM	13	23.6
	HT-DM-CVS-HL	8	14.5
	None	13	23.6
Dominant hand	Right	53	96.4
	Left	2	3.6
Stroke type	Hemorrhagic	7	12.7
	Ischemic	48	87.3
Involved side	Right	26	47.3
	Left	29	52.7
Lesion localization	TACI	14	25.5
	PACI	23	41.8
	POCI	18	32.7
Repeat stroke history	Yes	15	27.3
	No	40	72.7
Intensive care history	Yes	34	61.8
	No	21	38.2

HT: Hypertension; DM: Diabetes mellitus; CVS: Cardiovascular disease; HL: Hyperlipidemia; TACI: Total anterior circulatory infarcts; PACI: Partial anterior circulatory infarcts; POCI: Posterior circulatory infarcts.

The EAT-10 T1 score did not change statistically in T2 ($p=1.0$), and it decreased statistically in T3 ($p<0.001$).

Aspiration pneumonia (AP) was seen in only 4 (7.3%) of the patients during the follow-ups, but none of the patients died. The mean GUSS T1 score of the patients who developed AP during the follow-ups was 9.25 ± 1.89 , while the mean GUSS T1 score of the patients who did not develop AP was 13.96 ± 4.4 . Patients who developed AP were in the VFS group and aspiration was detected in these patients.

EVALUATION OF CLINICAL PARAMETERS ACCORDING TO GUSS AND EAT-10 T1 SCORES

The relationship between reflex, oral range of motion, oxygen demand, and incontinence characteris-




FIGURE 2: Algorithm of patients who underwent VFS according to GUSS T1 result. VFS: Videofluoroscopy; GUSS: Gugging Swallowing Screen.

tics of the patients with GUSS and EAT-10 T1 scores is shown in Table 2.

GUSS T1 score and BMI score were found to have a statistically significant positive correlation ($p<0.05$). There was no statistically significant difference between patients' gender, smoking status, alcohol use status, and comorbidity characteristics and GUSS and EAT-10 T1 scores ($p>0.05$). The correlation of GUSS and EAT-10 T1 score with clinical parameters is shown in Table 3.

EVALUATION OF CLINICAL PARAMETERS ACCORDING TO GUSS AND EAT-10 SCORE CHANGES AFTER THE REHABILITATION

The relationship between the oral hygiene and stroke characteristics of the patients and the GUSS and EAT-10 score changes after the rehabilitation is shown in Table 4.

There was no statistically significant difference between the patients' tongue and lip range of motion, palatal and gag reflexes, speech disorder, and lesion localization and the change in GUSS and EAT-10 scores after the rehabilitation ($p>0.05$).

DISCUSSION

Dysphagia is a significant cause of mortality and morbidity in stroke patients. Swallowing evaluation

TABLE 2: Distribution of GUSS and EAT-10 T1 scores according to reflex, oral range of motion, oxygen demand and incontinence characteristics of the patients.

Variables		GUSS T1 score		EAT-10 T1 score	
		($\bar{X}\pm$ SD)	p value	($\bar{X}\pm$ SD)	p value
Gag reflex	Yes	15.03±3.91	<0.001*	16.41±10.14	0.044*
	No	10.19±3.76		22.25±7.69	
Tongue movements	Full	14.24±4.13	0.003*	16.96±9.54	0.010*
	Limited	8.50±3.62		27.50±6.50	
Lip movements	Full	14.37±4.07	0.001*	16.90±9.63	0.013*
	Limited	8.43±3.31		26.43±6.58	
Palatal movements	Symmetrical	14.39±4.02	0.006*	16.65±8.99	0.007*
	Asymmetrical	9.67±4.53		25.56±10.85	
Aphasia	No	14.73±3.89	0.001*	16.24±9.25	0.013*
	Yes	10.36±4.41		23.57±9.59	
Oxygen requirement	Yes	10.7±4.92	0.028*	16.41±10.14	0.005*
	No	14.27±4.09		22.25±7.69	
Incontinence	Yes	11.65±4.16	0.013*	19.5±8.09	0.528
	No	14.74±4.23		17.31±10.67	

*p<0.05; GUSS: Gugging Swallow Screen; EAT-10: Eating Assessment Tool-10; SD: Standard deviation.

TABLE 3: Correlation of clinical parameters with GUSS and EAT-10 T1 score.

Variables	GUSS before rehabilitation program		EAT-10 before rehabilitation program	
	r	p value	r	p value
NIHSS	-0.383	0.004*	0.257	0.058
NIHSS facial paralysis	-0.079	0.566	-0.055	0.690
NIHSS aphasia	-0.411	0.002**	0.313	0.020*
NIHSS dysarthria	-0.237	0.082	0.090	0.513
BI	0.491	<0.001*	-0.323	0.016*
FAC	0.348	0.009*	-0.191	0.163

*p<0.05; GUSS: Gugging Swallow Screen; EAT-10: Eating Assessment Tool-10; NIHSS: National Institutes of Health Stroke Scale; BI: Barthel Index; FAC: Functional Ambulation Classification; r: Correlation coefficient.

TABLE 4: Distribution of GUSS and EAT-10 score changes after rehabilitation protocol according to oral hygiene and stroke characteristics of patients.

Variables		GUSS score change		EAT-10 score change	
		($\bar{X}\pm$ SD)	p value	($\bar{X}\pm$ SD)	p value
Oral hygiene	Poor	2.42±2.87	0.005*	8.33±7.35	0.984
	Moderate	2.22±2.66		9.00±8.11	
	Good	4.80±2.98		8.20±6.12	
Stroke type	Hemorrhagic	2.29±2.21	0.378	2.14±3.72	0.002*
	Ischemic	3.33±3.13		9.5±7.07	
History of recurrent stroke	No	3.95±3.2	0.002*	8.6±7.28	0.977
	Yes	1.2±1.01		8.47±7.03	

*p<0.05; GUSS: Gugging Swallow Screen; EAT-10: Eating Assessment Tool-10; SD: Standard deviation.

of patients should be performed starting from the acute period and patients with dysphagia should be followed up by giving a rehabilitation protocol. In

this study, acute stroke patients with dysphagia were given an individual rehabilitation protocol and followed up.

In the literature, it has been stated that early dysphagia screening with a standard screening protocol reduces the incidence of dysphagia-associated pneumonia and that screening methods and advanced examination methods should be used effectively and accurately.¹⁵ It is more appropriate to start the dysphagia screening test with a non-liquid bolus, as most acute stroke patients have difficulty swallowing liquid boluses and safer swallowing is observed in boluses with increased viscosity such as nectar and pudding. In this study, the GUSS test was applied to all patients because it allows the use of multiple consistencies, starting with pudding, allowing gradual evaluation, and at the same time providing dietary recommendations to minimize the risk of aspiration during administration.¹⁶

Recent studies have suggested that early dysphagia rehabilitation may improve outcomes for severe AP.¹⁷ Stegemann et al. stated that more than 90% of patients staying in the long-term care center had comorbidities requiring the use of solid form drugs and had serious difficulties in swallowing solid form drugs, and they stated that it would be easier to maintain a safe oral administration in the treatment of comorbidity with early dysphagia rehabilitation.¹⁸

In a study in which an exercise protocol was created for patients with dysphagia, patients were given a standard exercise program in the form of a home program once a week for 8 weeks under the supervision of a therapist. Improvement in dysphagia was observed in the follow-ups with VFS.¹⁹ In the present study, the patients were treated with an individual rehabilitation program, not a standard rehabilitation program, and a significant improvement was found in the post-treatment dysphagia scores of the patients, similar to previous studies.^{6,17}

The occurrence of pneumonia after stroke was shown to increase mortality and overall hospital care costs 3 times.²⁰ In a study that followed patients with dysphagia for 3 months after acute stroke, pneumonia was detected in 12% of the patients, while mortality was observed in 20%.²¹ In this study, AP developed in only 4 patients (7.3%) during follow-up and none of the patients died. The reason why this rate was less than the ones reported in the literature maybe that the rehabilitation program for dysphagia was applied while following the patients.

While there are many prospective studies evaluating the risk of aspiration after stroke, there exist only a few studies investigating predictors of dysphagia.^{2,22} In studies comparing stroke patients with and without dysphagia to determine predictors of dysphagia, no significant difference was found in terms of age, gender, smoking, alcohol use, and systemic comorbidities, consistent with this study.^{1,23,24}

In a study, no definite relationship was found between dysphagia and oxygen demand.¹ In this study, it was observed that the severity of dysphagia increased in patients who needed oxygen. Coordinated swallowing and respiration support this result. Neurological diseases accompanied by respiratory dysfunction may cause dysphagia and may be a risk factor for aspiration.²⁵

The importance of prolonged dysphagia and dysphagia prognosis is emphasized, as dysphagia may prolong and even become permanent in patients after acute stroke, affecting quality of life. In a study evaluating the long-term predictors of dysphagia in patients with acute stroke, some parameters such as understanding, expression, palatal movements, gag reflex, and voluntary cough were found to be significantly different between the transient and prolonged dysphagia group.²⁴ In this study, the severity of dysphagia was found to be high in patients with limited tongue and lip movement, palatal asymmetry, weak/no gag reflex, and patients with aphasia. It is thought that these parameters, which are easily evaluated in clinical practice in a short time and cost-free, will contribute to the determination of dysphagia severity and aspiration, and will help to arrange a suitable rehabilitation protocol for the patient.

The NIHSS is widely used to determine the general severity of neurological impairment in acute stroke. There exist several studies investigating the relationship between stroke severity and dysphagia. Studies have shown that stroke patients with dysphagia have a higher NIHSS score, which highlights the importance of NIHSS, an easy and quick test compared to costly clinical swallowing protocols that take time and require equipment. Studies investigating the relationship between stroke severity and dysphagia

have shown that stroke patients with dysphagia have a higher NIHSS score, similar to this study.^{2,22,26}

In this study, it was determined that there was a statistically significant positive correlation between the dysphagia score and the Barthel score. Previous studies also support this result. It is observed that the level of dependence in daily living activities of stroke patients negatively affects dysphagia.^{1,24,27}

The number of functional teeth is accepted as a very important parameter in the evaluation of oral health.²⁸ In the literature, a holistic evaluation of oral structures is recommended. Poor oral hygiene was found to have high specificity but low sensitivity in detecting aspiration in the post-stroke population.^{29,30} In this study, it was determined that the swallowing functions of patients with good oral hygiene improved significantly compared to those with poor oral hygiene. This shows the importance of explaining, demonstrating, and maintaining oral hygiene to the patient in preventing AP and gaining swallowing function.

Consistent with this study, it was shown in a study that whether the stroke type is hemorrhagic or ischemic does not affect the severity of dysphagia; however, there are limited studies in the related literature. Paciaroni et al. observed that the frequency of dysphagia increased in patients with hemorrhagic stroke compared to ischemic stroke, and they thought that this might be related to the severity of stroke.^{1,31} In another study, dysphagia was shown more frequently in hemorrhagic patients and it was reported that hemorrhagic stroke may be an independent predictor of prolonged dysphagia.³² In this study, a significant difference was found between the changes in EAT-10 score of patients with ischemic and hemorrhagic stroke before and after the rehabilitation protocol application. Ischemic stroke patients had a better prognosis for recovery and therefore dysphagia, which suggests that it is important to follow up patients with hemorrhagic stroke for a longer period of time. There is a need for more comprehensive studies showing the relationship between hemorrhagic stroke and dysphagia.

No comprehensive dysphagia screening and rehabilitation protocol was found in the literature, as in this study, which screened acute stroke patients for

dysphagia, followed the patient with an individual rehabilitation protocol in patients with dysphagia and evaluated the clinical parameters that could affect the severity and prognosis of dysphagia. Dysphagia screening protocols have been established for early detection of dysphagia in studies, and in a small number of studies, standard treatment protocols have been established.^{19,33-35} In our study, we aimed to emphasize individual rehabilitation and reduce morbidity due to dysphagia by presenting the rehabilitation protocol and follow-up results in patients with dysphagia detected by evaluating swallowing in stroke patients from the acute period to the literature.

The most important and also strong aspect of the study is that dysphagia evaluation was performed in the acute period. Thus, an individual rehabilitation was arranged by making dietary modifications for the patients without the development of possible AP, dehydration, and malnutrition. Dysphagia, which is an important cause of mortality and morbidity, was evaluated in many aspects by following the patients closely, taking into account the severity of dysphagia and other factors affecting its prognosis. It is thought that this study will contribute to the literature by following up the patients by applying early screening and early dysphagia rehabilitation protocol and presenting the effects of the results on clinical parameters.

The absence of a control group in the early dysphagia rehabilitation can be cited as one of the limitations of the study. However, it is thought that leaving the control group without treatment would not comply with ethical principles. Another limitation is that the patients were not followed up for more than one month after discharge in terms of mortality and morbidity. The results need to be supported by studies with a higher number of patients and with long-term follow-up with controls.

CONCLUSION

In this study, it was observed that close monitoring of acute stroke patients with dysphagia with an individual rehabilitation protocol improved dysphagia scores and reduced morbidity that may develop in patients. Dysphagia, which is an important cause of mortality and morbidity in stroke patients, should be

evaluated in the acute period and followed with an individual rehabilitation protocol appropriate for the patient.

Acknowledgements

The authors would like to thank all patients for their support.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct con-

nection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

REFERENCES

1. Baroni AF, Fábio SR, Dantas RO. Risk factors for swallowing dysfunction in stroke patients. *Arq Gastroenterol.* 2012;49:118-24. PMID: 22766998.
2. Henke C, Foerch C, Lapa S. Early screening parameters for dysphagia in acute ischemic stroke. *Cerebrovasc Dis.* 2017;44:285-90. PMID: 28903096.
3. Jauch EC, Saver JL, Adams HP Jr, et al; American Heart Association Stroke Council; Council on Cardiovascular Nursing; Council on Peripheral Vascular Disease; Council on Clinical Cardiology. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2013;44:870-947. PMID: 23370205.
4. Casaubon LK, Boulanger JM, Blacquiére D, et al; Heart and Stroke Foundation of Canada Canadian Stroke Best Practices Advisory Committee. Canadian Stroke Best Practice Recommendations: Hyperacute Stroke Care Guidelines, Update 2015. *Int J Stroke.* 2015;10:924-40. PMID: 26148019.
5. Moraes DP, Sassi FC, Mangilli LD, et al. Clinical prognostic indicators of dysphagia following prolonged orotracheal intubation in ICU patients. *Crit Care.* 2013;17:R243. PMID: 24138781; PMCID: PMC4056041.
6. Krajczyk E, Krajczyk M, Luniewski J, et al. Assessment of the effects of dysphagia therapy in patients in the early post-stroke period: a randomised controlled trial. *Neurol Neurochir Pol.* 2019;53:428-34. PMID: 31724147.
7. Umay EK, Gürçay E, Bahçeci K, et al. Validity and reliability of Turkish version of the gugging swallowing screen test in the early period of hemispheric stroke. *Neurol Sci Neurophysiol* 2018;35:6-13. doi: 10.5152/NSN.2018.10436
8. Belafsky PC, Mouadeb DA, Rees CJ, et al. Validity and reliability of the Eating Assessment Tool (EAT-10). *Ann Otol Rhinol Laryngol.* 2008;117:919-24. PMID: 19140539.
9. Demir N, Serel Arslan S, İnal Ö, et al. Reliability and validity of the Turkish Eating Assessment Tool (T-EAT-10). *Dysphagia.* 2016;31:644-9. PMID: 27405421.
10. Karaduman A, Serel S, Ünlüer Ö ve ark. [The Penetration Aspiration Scale: an interrater reliability study]. *Fizyoter Rehabil* 2012;23:151-5. <https://dergi-park.org.tr/en/download/article-file/138070>
11. Schlegel D, Kolb SJ, Luciano JM, et al. Utility of the NIH Stroke Scale as a predictor of hospital disposition. *Stroke.* 2003;34:134-7. PMID: 12511764.
12. Küçükdeveci AA, Yavuzer G, Tennant A, et al. Adaptation of the modified Barthel Index for use in physical medicine and rehabilitation in Turkey. *Scand J Rehabil Med.* 2000;32:87-92. PMID: 10853723.
13. Holden MK, Gill KM, Magliozzi MR, et al. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther.* 1984;64:35-40. PMID: 6691052.
14. Ong JJ, Steele CM, Duizer LM. Sensory characteristics of liquids thickened with commercial thickeners to levels specified in the International Dysphagia Diet Standardization Initiative (IDDSI) framework. *Food Hydrocoll.* 2018;79:208-17. PMID: 29795963; PMCID: PMC5962271.
15. Eltringham SA, Kilner K, Gee M, et al. Impact of dysphagia assessment and management on risk of stroke-associated pneumonia: a systematic review. *Cerebrovasc Dis.* 2018;46:99-107. PMID: 30199856.
16. Warnecke T, Im S, Kaiser C, et al. Aspiration and dysphagia screening in acute stroke - the Gugging Swallowing Screen revisited. *Eur J Neurol.* 2017;24:594-601. PMID: 28322006.
17. Nakamura T, Kurosaki S. Effects of early dysphagia rehabilitation by speech-language-hearing therapists on patients with severe aspiration pneumonia. *Prog Rehabil Med.* 2020;5:20200020. PMID: 32908953; PMCID: PMC7471375.
18. Stegemann S, Gosch M, Breitkreutz J. Swallowing dysfunction and dysphagia is an unrecognized challenge for oral drug therapy. *Int J Pharm.* 2012;430:197-206. PMID: 22525080.
19. Balou M, Herzberg EG, Kamelhar D, et al. An intensive swallowing exercise protocol for improving swallowing physiology in older adults with radiographically confirmed dysphagia. *Clin Interv Aging.* 2019;14:283-8. PMID: 30804667; PMCID: PMC6375531.
20. Papavasileiou V, Milionis H, Smith CJ, et al. External validation of the pre-stroke independence, sex, age, National Institutes of Health Stroke Scale (ISAN) score for predicting stroke-associated pneumonia in the athens stroke registry. *J Stroke Cerebrovasc Dis.* 2015;24:2619-24. PMID: 26347399.
21. Fernández-Pombo A, Seijo-Raposo IM, López-Osorio N, et al. Lesion location and other predictive factors of dysphagia and its complications in acute stroke. *Clin Nutr ESPEN.* 2019;33:178-82. PMID: 31451257.
22. Joundi RA, Martino R, Saposnik G, et al. Predictors and outcomes of dysphagia screening after acute ischemic stroke. *Stroke.* 2017;48:900-6. PMID: 28275200.
23. Leite KKA, Sassi FC, Medeiros GC, et al. Clinical swallowing prognostic indicators in patients with acute ischemic stroke. *Arq Neuropsiquiatr.* 2019;77:501-8. PMID: 31365642.
24. Broadley S, Croser D, Cottrell J, et al. Predictors of prolonged dysphagia following acute stroke. *J Clin Neurosci.* 2003;10:300-5. PMID: 12763332.
25. Nilsson H, Ekberg O, Olsson R, et al. Dysphagia in stroke: a prospective study of quantitative aspects of swallowing in dysphagic patients. *Dysphagia.* 1998;13:32-8. PMID: 9391228.
26. Jeyaseelan RD, Vargo MM, Chae J. National Institutes of Health Stroke Scale (NIHSS) as an early predictor of poststroke dysphagia. *PM R.* 2015;7:593-8. PMID: 25556573.
27. Broadley S, Cheek A, Salonikis S, et al. Predicting prolonged dysphagia in acute stroke: the Royal Adelaide Prognostic Index for Dysphagic Stroke (RAPIDS). *Dysphagia.* 2005;20:303-10. PMID: 16633876.

-
28. Fukai K, Takiguchi T, Ando Y, et al. Critical tooth number without subjective dysphagia. *Geriatr Gerontol Int.* 2011;11:482-7. PMID: 21702874.
 29. McCullough GH, Rosenbek JC, Wertz RT, et al. Utility of clinical swallowing examination measures for detecting aspiration post-stroke. *J Speech Lang Hear Res.* 2005;48:1280-93. PMID: 16478371.
 30. Felix CC, Joseph ME, Daniels SK. Clinical decision making in patients with stroke-related dysphagia. *Semin Speech Lang.* 2019;40:188-202. PMID: 31158903.
 31. Paciaroni M, Mazzotta G, Corea F, et al. Dysphagia following Stroke. *Eur Neurol.* 2004;51:162-7. PMID: 15073441.
 32. Toscano M, Cecconi E, Capiluppi E, et al. Neuroanatomical, Clinical and cognitive correlates of post-stroke dysphagia. *Eur Neurol.* 2015;74:171-7. PMID: 26492033.
 33. Taveira I, Silva S, Bonaça Í, et al. Recognizing dysphagia: implementation of an in-hospital screening protocol. *Ir J Med Sci.* 2021;190:605-8. PMID: 32959220.
 34. Sherman V, Flowers H, Kapral MK, et al. Screening for dysphagia in adult patients with stroke: assessing the accuracy of informal detection. *Dysphagia.* 2018;33:662-9. PMID: 29497830.
 35. Perry SE, Miles A, Fink JN, et al. The dysphagia in stroke protocol reduces aspiration pneumonia in patients with dysphagia following acute stroke: a clinical audit. *Transl Stroke Res.* 2019;10:36-43. PMID: 29603057.

ARTICLE IN PRESS